

Machine Learning Theory (CS 6783)

Lecture 16: Supplement

1 Two Equivalent Definitions of Convexity

For this section, say \mathcal{B} is some vector space equipped with norm $\|\cdot\|$ and \mathcal{B}^* be the dual space equipped with dual norm $\|\cdot\|_*$. For simplicity think of \mathcal{B} and \mathcal{B}^* to simply be \mathbb{R}^d . The following are two equivalent definitions of convex functions.

Definition 1. A function $f : \mathcal{B} \mapsto \mathbb{R}$ is said to be convex if for all $x, y \in \mathcal{B}$ and any $\alpha \in [0, 1]$,

$$f(\alpha x + (1 - \alpha)y) \leq \alpha f(x) + (1 - \alpha)f(y)$$

The following is an equivalent definition in terms of gradients.

Definition 2. A function $f : \mathcal{B} \mapsto \mathbb{R}$ is said to be convex if for all $x, y \in \mathcal{B}$

$$f(x) \leq f(y) + \langle \nabla f(x), x - y \rangle$$

Why are the two definitions equivalent?

(**1** \Rightarrow **2**) First lets show that the first definition implies the second. Note that by definition of directional derivative:

$$\begin{aligned} \langle \nabla f(x), y - x \rangle &= \lim_{\alpha \rightarrow 0} \frac{f(x + \alpha(y - x)) - f(x)}{\alpha} \\ &= \lim_{\alpha \rightarrow 0} \frac{f((1 - \alpha)x + \alpha y) - f(x)}{\alpha} \\ &\leq \lim_{\alpha \rightarrow 0} \frac{(1 - \alpha)f(x) + \alpha f(y) - f(x)}{\alpha} \\ &= \lim_{\alpha \rightarrow 0} \frac{\alpha(f(y) - f(x))}{\alpha} \\ &= f(y) - f(x) \end{aligned}$$

Rearranging we see that definition 1 implies definition 2.

(**2** \Rightarrow **1**) Now to prove the other direction, starting with definition 2, we have the following two inequalities:

$$\begin{aligned} f(\alpha x + (1 - \alpha)y) &\leq f(x) + \langle \nabla f(\alpha x + (1 - \alpha)y), \alpha x + (1 - \alpha)y - x \rangle \\ &= f(x) + (1 - \alpha) \langle \nabla f(\alpha x + (1 - \alpha)y), y - x \rangle \end{aligned}$$

Similarly,

$$\begin{aligned} f(\alpha x + (1 - \alpha)y) &\leq f(y) + \langle \nabla f(\alpha x + (1 - \alpha)y), \alpha x + (1 - \alpha)y - y \rangle \\ &= f(y) + \alpha \langle \nabla f(\alpha x + (1 - \alpha)y), y - x \rangle \end{aligned}$$

Hence summing up α times the first inequality and $1 - \alpha$ times the second we end up with

$$\begin{aligned} f(\alpha x + (1 - \alpha)y) &\leq \alpha f(x) + (1 - \alpha)f(y) + (1 - \alpha)\alpha \langle \nabla f(\alpha x + (1 - \alpha)y), y - x \rangle + \\ \alpha(1 - \alpha) \langle \nabla f(\alpha x + (1 - \alpha)y), x - y \rangle & \\ &= \alpha f(x) + (1 - \alpha)f(y) \end{aligned}$$

This completes the proof.